THATCH BUSTER

SOIL



BENEFITS:

ACCELERATES DECOMPOSITION OF CELLULOSE DECREASES INCIDENCE OF UNHEALTHY THATCH LAYERING PROMOTES CONVERSION OF DEAD CLIPPINGS TO USABLE NUTRIENTS

Thatch Buster is a concentrated cellulase enzyme product that duplicates those in nature, breaking down the major component of dead plant tissue.



FLORATINE Rooted in Science

THATCH BUSTER

UNRAVEL CELLULOSE

Maintenance of dormant enzymes in solution requires special measures, as does initiation of their activity. When activated, precautions must be taken to avoid mixing with harsh chemicals. The enzymes in Thatch Buster are more active at higher temperatures. Do not apply when soil temperatures are at 60°F (15°C) or lower. Treated areas should be kept moist to ensure enzymatic activity.

PROFESSIONAL USE GUIDELINES (5)



APPLICATION:

Best results will be realized when ambient air temperatures are greater than 70°F (21°C). Material should be watered in to the interface of soil and thatch. Where thatch is very dense, *Pervade* can be added to the tank mix to accelerate penetration.

RATE OF APPLICATION:

Apply 3 fl oz / 1000 sq ft (10 L/ha). Apply every 3-4 weeks for maintenance applications.

COMPATIBILITY:

Do not mix with non-Floratine adjuvants as they may damage the enzymes.

Net Weight: 8.18 lb/gal (0.980 kg/L)

Floratine Products Group, Inc. 355 East South Street • Collierville, TN 38017 901.853.2898 • www.floratine.com



FLORATINE Rooted in Science